

INFUSION & MEDICAL CENTER

| Pa | itient Name | DOB | Patient Phone/Cell # |
|-----|---|--|---|
| | Patient demographic and insurance in | formation to be faxed | d with Infusion Order Form |
| M | edical Information (Please complete/select | appropriate diagns | osis): |
| | Primary Diagnosis:Relapsing Multiple S | clerosis | ICD-10 Code: G35 |
| | Allergies: | | (or attach list) |
| Cli | inical Information – Please fax with Infusio | n Order Form: | Patient |
| | • Clinical notes & labs supporting primary diagno | osis | Weight: lbs. |
| | Hepatitis B Screening Results | | Height: in. |
| _ | OCREVILO | ® (ocrelizumab) | J Code: J2350 |
| | OCREVUS | • | |
| Dı | rug Order: | | |
| Dı | rug Order: | d into 2 infusions | |
| | rug Order: Loading Dose: Ocrevus 600 mg IV divide Administer 300 mg IV over 2.5 hours on 0 week | d into 2 infusions and 2 weeks. | # Refills (Recommend 1 Refills) |
| | rug Order: Loading Dose: Ocrevus 600 mg IV divide Administer 300 mg IV over 2.5 hours on 0 week | d into 2 infusions and 2 weeks. every 6 months | |
| | rug Order: Loading Dose: Ocrevus 600 mg IV divide Administer 300 mg IV over 2.5 hours on 0 week Maintenance Dose: Ocrevus 600 mg IV Administer 600 mg IV over 2 hours or as tolerate Pre-Medication Orders: | d into 2 infusions and 2 weeks. every 6 months ed – 6 months after the | most recent infusion IV, and methylprednisolone 125 mg IV |
| | rug Order: Loading Dose: Ocrevus 600 mg IV divide Administer 300 mg IV over 2.5 hours on 0 week Maintenance Dose: Ocrevus 600 mg IV Administer 600 mg IV over 2 hours or as tolerate Pre-Medication Orders: Acetaminophen 650 mg PO, Di | d into 2 infusions and 2 weeks. every 6 months ed – 6 months after the phenhydramine 50 mg fusion *Adjust | most recent infusion IV, and methylprednisolone 125 mg IV to patient's needs |
| | Loading Dose: Ocrevus 600 mg IV divide Administer 300 mg IV over 2.5 hours on 0 week Maintenance Dose: Ocrevus 600 mg IV Administer 600 mg IV over 2 hours or as tolerate Pre-Medication Orders: Acetaminophen 650 mg PO, Di Administered 30 min prior to in | d into 2 infusions and 2 weeks. every 6 months ed – 6 months after the open shadow and shadow and shadow are shadow as a sh | most recent infusion IV, and methylprednisolone 125 mg IV to patient's needs |
| | Loading Dose: Ocrevus 600 mg IV divide Administer 300 mg IV over 2.5 hours on 0 week Maintenance Dose: Ocrevus 600 mg IV Administer 600 mg IV over 2 hours or as tolerate Pre-Medication Orders: Acetaminophen 650 mg PO, Di Administered 30 min prior to in Famotidine 20 mg administered IV 30 minutes p | d into 2 infusions and 2 weeks. every 6 months ed – 6 months after the phenhydramine 50 mg ifusion *Adjust | most recent infusion IV, and methylprednisolone 125 mg IV to patient's needs infusion |
| | Loading Dose: Ocrevus 600 mg IV divide Administer 300 mg IV over 2.5 hours on 0 week Maintenance Dose: Ocrevus 600 mg IV Administer 600 mg IV over 2 hours or as tolerate Pre-Medication Orders: Acetaminophen 650 mg PO, Di Administered 30 min prior to in Famotidine 20 mg administered IV 30 minutes go | d into 2 infusions and 2 weeks. every 6 months ed – 6 months after the phenhydramine 50 mg ifusion *Adjust orior to the start of the adverse reaction that hese services, I am auti | IV, and methylprednisolone 125 mg IV to patient's needs infusion may occur per approved ADR Protocol. norizing Intramed Plus |
| | Loading Dose: Ocrevus 600 mg IV divide Administer 300 mg IV over 2.5 hours on 0 week Maintenance Dose: Ocrevus 600 mg IV over 2 hours or as tolerate Pre-Medication Orders: Acetaminophen 650 mg PO, Dig Administered 30 min prior to in Famotidine 20 mg administered IV 30 minutes potential. Other: Adverse Drug Reaction Protocol: Manage any By signing this form and utilizing to serve as my prior authorization agen | d into 2 infusions and 2 weeks. every 6 months ed – 6 months after the phenhydramine 50 mg ifusion *Adjust orior to the start of the adverse reaction that these services, I am autil t with medical and pha | IV, and methylprednisolone 125 mg IV to patient's needs infusion may occur per approved ADR Protocol. norizing Intramed Plus rmacy insurance providers. |
| | Loading Dose: Ocrevus 600 mg IV divide Administer 300 mg IV over 2.5 hours on 0 week Maintenance Dose: Ocrevus 600 mg IV over 2 hours or as tolerate Pre-Medication Orders: Acetaminophen 650 mg PO, Di Administered 30 min prior to in Famotidine 20 mg administered IV 30 minutes potential Other: Adverse Drug Reaction Protocol: Manage any By signing this form and utilizing to | d into 2 infusions and 2 weeks. every 6 months ed – 6 months after the phenhydramine 50 mg ifusion *Adjust orior to the start of the adverse reaction that hese services, I am autil t with medical and pha | IV, and methylprednisolone 125 mg IV to patient's needs infusion may occur per approved ADR Protocol. norizing Intramed Plus rmacy insurance providers. Date: Date: n permitted |

FAX ALL INFORMATION CENTRAL FAX 803.999.1754

INFUSION CENTER LOCATIONS

BERKELEY CHARLESTON COLUMBIA GREENVILLE CENTRAL INTAKE PHONE 803.999.1760