

## **INFUSION & MEDICAL CENTER**

| 1.Patient Name                              |   | DOB                 | Patient Phone/Cell #  |
|---|---|---------------------|-----------------------|
| Patient de                                  | mographic and insurance information                                 | n to be faxed wit   | h Infusion Order Form |
| 2.Medical Informat                          | ion (Please select primary diagr                                    | osis and com        | plete ICD-10 Code):   |
| Primary Diagnosis:                          | Primary Hyperoxaluria Type 1  | ICD-10 Code: E72.53 |                       |
|   | Other:  | IC                  | D-10 Code:            |
| Allergies:                                  |   |                     | (or attach list)      |
|   | on — Please fax with Infusion O<br>ion supporting primary diagnosis | rder Form:          |                       |
| Recent Lab/Test Results including:          |   |                     | Patient               |
| o AGXT genetic test<br>o Urine or plasma ox |   |                     | Weight:lb             |
| Medication List                             |   |                     | Height in             |
| <ul> <li>Continuation of ther</li> </ul>    | apy: Oxlumo start date:   |                     |                       |
| o Attach notes show                         | ring a reduction in urinary or plasma oxa                           | late levels compa   | red to baseline       |

## 4. Drug Order:

# OXLUMO<sup>®</sup> (Lumasiran)

J Code: J0224

**OXLUMO**®

### New Start / Loading Dose

| Patient Weight     | Dose    | Directions                         | Doses/Refills               |
|--------------------|---------|------------------------------------|-----------------------------|
| Less than 10 kg    | 6 mg/kg |                                    |                             |
| □ 10 kg to < 20 kg | 6 mg/kg | Inject subcutaneously once monthly | Doses authorized: 3 (three) |
| 🖵 20 kg and above  | 3 mg/kg |                                    |                             |

**Maintenance Regimen** (to be initiated 1 month following the final administration of the loading dose):

| Patient Weight     | Dose    | Directions Doses/Refills               |                           |
|--------------------|---------|--|---------------------------|
| 🖵 Less than 10 kg  | 3 kg/kg | Inject subcutaneously once monthly     | Refills: (Recommend 11)   |
| □ 10 kg to < 20 kg | 6 mg/kg | In is staub suten sough, such 2 months | Defile: (Decement en d.2) |
| 20 kg and above    | 3 mg/kg | Inject subcutaneously every 3 months   | Refills: (Recommend 3)    |

#### **Pre-Medication Orders:**

No Pre-medications are recommended based on manufacturer guidelines.

Adverse Drug Reaction Protocol: Manage any adverse reaction that may occur per approved ADR Protocol.

By signing this form and utilizing our services, I am authorizing Intramed Plus to serve as my prior authorization agent with medical and pharmacy insurance providers.

| FAX ALL INFORMATION<br>CENTRAL FAX <b>803.999.1754</b> | CENTRAL INTAKE PHONE<br>803.999.1750 |  |  |
|--|--------------------------------------|--|--|
| Printed Physician's Name with Credentials:             | Phone #:                             |  |  |
| Dispense as written                                    | Substitution permitted               |  |  |
| 5. Physician Signature:                                | / Date:                              |  |  |